

ON THE ONLY THREE SHORT-DISTANCE STRUCTURES WHICH CAN BE DESCRIBED BY LINEAR OPERATORS

Achim Kempf

Department of Applied Mathematics and Theoretical Physics
and Corpus Christi College, Cambridge University, United Kingdom

Abstract

We point out that if spatial information is encoded through linear operators X_i , or ‘infinite-dimensional matrices’ with an involution $X_i^* = X_i$ then these X_i can only describe either continuous, discrete or certain “fuzzy” space-time structures. We argue that the fuzzy space structure may be relevant at the Planck scale. The possibility of this fuzzy space-time structure is related to subtle features of infinite dimensional matrices which do not have an analogue in finite dimensions. For example, there is a slightly weaker version of self-adjointness: symmetry, and there is a slightly weaker version of unitarity: isometry. Related to this, we also speculate that the presence of horizons may lead to merely isometric rather than unitary time evolution.

1. NONCOMMUTING LIMITS AND INFINITE-DIMENSIONAL MATRICES

Since limits do not have to be interchangeable, infinite-dimensional matrices possess subtle features without analogues among finite dimensional matrices. Let us first look at the trivial example

$$k(n_1, n_2, a) = \frac{n_1 f(a) + n_2 g(a)}{n_1 + n_2}, \quad n_1, n_2 \in \mathbb{N} \quad (1.1)$$

where f and g are regular functions. Now $\lim_{n_1 \rightarrow \infty} \lim_{n_2 \rightarrow \infty} k(n_1, n_2, a) = g(a)$. It seems that when n_1 and n_2 are taken to infinity the information on f is lost. Of course it is only hidden and is not lost, because we can find f at infinity by approaching infinity on other paths, e.g. $\lim_{n_2 \rightarrow \infty} \lim_{n_1 \rightarrow \infty} k(n_1, n_2, a) = f(a)$. In this sense, infinity is able to store away things in such a way that various information can be retrieved by checking at various corners at infinity.

Now take an infinite dimensional matrix X_{ij} , ($i, j = 1, 2, \dots, \infty$) which obeys $X_{ij} = X_{ji}^*$. On the set D of all vectors which have only a finite (but arbitrarily large) number of nonzero entries, X is clearly symmetric, i.e. all its expectation values are real:

$$\langle v | X | v \rangle = \sum_{n_1, n_2=1}^{\infty} v_{n_1}^* X_{n_1 n_2} v_{n_2} \in \mathbb{R} \quad \forall v \in D \quad (1.2)$$

To take matrix elements of X between *generic* vectors is, however, nontrivial: The scalar product of $|v\rangle$ with $|X.w\rangle$ does not necessarily coincide with the scalar product of $|X.v\rangle$ with $|w\rangle$, because for

$$k(v, w, n_1, n_2) = \sum_{i=1}^{n_1} \sum_{j=1}^{n_2} v_i^* X_{ij} w_j \quad (1.3)$$

the two limits $n_1 \rightarrow \infty$ and $n_2 \rightarrow \infty$ have no reason to commute, i.e. in general we have

$$\lim_{n_1 \rightarrow \infty} \lim_{n_2 \rightarrow \infty} k(v, w, n_1, n_2) \neq \lim_{n_2 \rightarrow \infty} \lim_{n_1 \rightarrow \infty} k(v, w, n_1, n_2) \quad (1.4)$$

even if both v and w are square summable¹. We are now aware that in this case, roughly speaking, some information contained in X may be somewhat nontrivially stored.

Let us ask whether such subtleties could play a role in whatever will eventually be the fundamental theory of quantum gravity.

2. GRAVITY AT THE PLANCK SCALE AND NONCOMMUTATIVE GEOMETRY

Gravity has its fingers at both ends of the length spectrum: the extrapolation of general relativity and quantum theory, as we know them, leads to apparent paradoxes both at small and at large scales. A truly fundamental theory of quantum gravity will have to resolve these problems and explain such tricky issues like the structure of space-time at the Planck scale or also the apparent contradiction of unitary time evolution with black hole information loss.

Instead of going into the details of any particular approach to quantum gravity, such as string theory, let us step back for a moment to look from some distance at this problem of unifying quantum theory and gravity. With the above issues in mind, can we see any assumptions that are conventionally taken for granted in quantum theory, which might prove to be relaxed in a fundamental theory of quantum gravity?

One suggestion, going back to the 1940s [1] is that spatial information will still be encoded through linear operators, say X_i , but that these may eventually prove to be noncommutative. The idea of noncommutative geometry has in the meanwhile led to the development of beautiful and powerful mathematics and has been supported by new theoretical evidence within various approaches to quantum gravity, see e.g. [2].

Let us now consider the possibility that the X_i , could also be non-self-adjoint, merely symmetric operators². Correspondingly, the suggestion is that certain transformations which one would normally expect to be unitary, may ultimately prove to be merely isometric. Let us look at the likely properties of an individual X_i more closely:

3. LINEAR X_i CAN ENCODE ONLY THREE DIFFERENT SHORT DISTANCE STRUCTURES

We expect that in whatever algebra the X_i may be found to live, the involution in the algebra acts as $X_i^* = X_i$. This is to ensure that all formal expectation values of

¹ Technically, this is of course the fact that a symmetric operator X may or may not have a self-adjoint extension and if it has, it may have a whole family of self-adjoint extensions. This will be addressed in detail below.

²This will also yield a possibility for a more subtle form of noncommutativity of the X_i , which is not seen on the algebraic level.

the X_i in Hilbert space representations are real. One might therefore be tempted to conclude that the X_i are self-adjoint operators in Hilbert space representations. Let us, however, recall the following crucial point: An infinite dimensional matrix X_i of which all expectation values are real is called symmetric, but it is not necessarily self-adjoint, as we just recalled above.

Generally, a matrix X which has exclusively real expectation values (i.e. which is by definition symmetric), describes one out of exactly three different spatial structures:

1. if symmetric and self-adjoint, X may have a continuous spectrum and thus describe a continuous space, or
2. if symmetric and self-adjoint, X may have a discrete spectrum, thus describing a lattice, or
3. if simple symmetric, X has no eigenvectors, and describes, as we will see, a "fuzzy" space.

These are the extreme possibilities and arbitrary mixtures of the three can occur: A general symmetric matrix can be self-adjoint on subspaces and a general self-adjoint matrix can have a mixed continuous and discrete (even fractal) spectrum.

Of course, this is a statement for generic symmetric operators which need not be represented as infinite dimensional matrices. But since we are dealing with separable Hilbert spaces, let us think of these operators in matrix representations. I hope this helps intuition by clearly keeping those infinities that are important here apart from the trivial infinitesimals and infinities of continuous representations which would be mere artifacts of having chosen a continuous representation.

4. HOW TO IDENTIFY THE THIRD SHORT DISTANCE STRUCTURE IN PRACTISE

Characteristic for the third type short distance structure is that a simple symmetric X does not have eigenvectors (if it had eigenvectors, it would be self-adjoint on the eigenspaces, but by definition simple symmetric operators are not self-adjoint even on subspaces). In order to identify the third short distance structure in practise, it would therefore be useful to have a functional which indicates the presence or absence of eigenvectors. Indeed, a suitable indicator functional is the smallest formal uncertainty in X , defined for any symmetric X on its dense domain D_X in a Hilbert space as:

$$(\Delta X)_0 := \inf_{\{v \in D_X | \langle v|v \rangle = 1\}} \langle v | (X - \langle v|X|v \rangle)^2 | v \rangle^{1/2} \quad (4.1)$$

The term $\langle v | (X - \langle v|X|v \rangle)^2 | v \rangle^{1/2}$ vanishes for eigenvectors, and only for eigenvectors. Thus, for types 1, 2 and the mixed cases, we have $(\Delta X)_0 = 0$. Alternatively, when the formal lower bound to spatial resolution $(\Delta X)_0$ is some value larger than zero, then we have that X is of the third type - describing an in this sense "fuzzy" geometry.

The stringy correction to the uncertainty relation (see e.g. [4]) provides an example:

$$\Delta x \Delta p \geq \frac{\hbar}{2} (1 + \beta (\Delta p)^2 + \dots) \quad (4.2)$$

As is easily verified, Eq.4.2 implies that the x -uncertainty is finitely bounded from below: $(\Delta X)_0 = \hbar \sqrt{\beta}$. Therefore, any operator X that obeys this uncertainty relation is of the third type, i.e. it is simple symmetric.

5. INDICATIONS THAT FUZZY GEOMETRIES ARE UV REGULAR AND COMPATIBLE WITH EXTERNAL SYMMETRIES

The ‘fuzzy’ geometries described by X_i which are of the third type appear to have attractive features.

As opposed to the second case, the lattice, the geometries of the third type may arise without breaking external symmetries: For example the uncertainty relation Eq.4.2 can be obtained from the commutation relation

$$[X, P] = i\hbar(1 + \beta P^2 + \dots) \quad (5.1)$$

for which

$$X \rightarrow X + c, \quad P \rightarrow P \quad (5.2)$$

is an algebra morphism. This proves that translation invariance *can* be preserved on third type geometries, even if, e.g., Eq.4.2 should ultimately arise from some more sophisticated fundamental theory.

Similarly, it has been shown that in more dimensions also rotation invariance can be preserved: One can find examples of X_i which obey translation and rotation invariant commutation relations that lead to uncertainty relations which imply that the X_i are of the third type (also, if the simple symmetric X_i commute, the form of the lowest order correction terms to the CCRs is unique with the fundamental length scale $(\Delta X)_0$ as the only free parameter), see [5].

As opposed to the first case, the continuum, the fuzzy geometries appear to also provide a natural ultraviolet cutoff:

The framework of quantum field theory may well only have limited validity, but it is at least an interesting indication that QFTs can be formulated on fuzzy geometries and are ultraviolet regular, see e.g. [6]: Technically, when Feynman rules are defined in position space the ultraviolet divergencies appear as the ill-definedness of products of propagators and vertices.

These products are ill-defined because they are then products of distributions like

$$\delta(x - x') = \langle x|x' \rangle \quad \text{and} \quad G(x, x') = \langle x|1/(\mathbf{p}^2 + m^2)|x' \rangle. \quad (5.3)$$

But why are δ and G distributions? They are distributions exactly because the vectors of maximal spatial localisation ‘ $|x\rangle$ ’ are non-normalisable for X_i which describe a continuum, i.e. which are of the type 1. To be precise: a vector of maximal localisation obeys $(\Delta X_i)_{|x\rangle} = (\Delta X)_0, \forall i$ and x denotes the formal expectation value $\langle x|X_i|x\rangle = x_i$. On the continuum they are of course the non-normalisable formal X_i -eigenfields.

Clearly, the vectors of maximal localisation are normalisable in the case that the X_i is of the second type, the lattice. But, it has also been shown that they are normalisable when the X_i are of the third type [7]. If the $|x\rangle$ are normalisable, δ and G are regular functions and their products are well-defined, from which follows that the UV divergencies are regularised.

While this shows that geometries described by X_i of the third type can preserve external symmetries and regularise the ultraviolet, we still have not picked up on our observation in the beginning that with these simple symmetric X_i , ‘some information may be stored somewhat nontrivially’.

6. THE DEGREES OF FREEDOM CUT OFF BEYOND THE PLANCK SCALE CAN REAPPEAR AS INTERNAL DEGREES OF FREEDOM

If a fundamental theory of quantum gravity does encode spatial information through X_i of the third type then those spatial degrees of freedom that are being cut off beyond the scale $(\Delta X)_0$ may reappear in a different form, as internal degrees of freedom:

Let us first look at QED. It can be formulated by specifying that the fields in the, say euclidean, field theoretical path integral are in a representation of the algebra $[P_i, P_j] = F_{ij}(X)$, $[X_i, P_j] = i\hbar\delta_{ij}$, $[X_i, X_j] = 0$. The gauge transformations are the group G of unitaries in the algebra of the X_i . We note that each unitary, say $e^{if(X)}$, can be written as a function of n elementary unitaries defined as

$$S_j = (X_j - i1)(X_j + i1)^{-1} \quad (6.1)$$

i.e. there always exists a g such that

$$e^{if(X)} = g(S_1, \dots, S_n) \quad (6.2)$$

To obtain a non-abelian gauge theory the isospinor structure would normally have to be introduced ‘by hand’.

Now if a fundamental quantum gravity theory does encode spatial information through fuzzy X_i then the group G of unitaries in the algebra of the X_i is generically non-abelian, even if the X_i are commutative:

The reason is that the S_i which arise from the X_i are merely isometric and require unitary extension. Generically, unitary extensions exist and differ by elements of some unitary group T (the unitary automorphisms of the deficiency spaces $L_i^+ = D_{S_i}^\perp$, $L_i^- = R_{S_i}^\perp$ of the X_i), see [8]. The resulting G therefore naturally splits into conventional unitaries which are functions of the S_i , and T . As will be shown in general and in detail in [9], at low energies the gauge transformations defined as the set of unitaries which commute with the X_i factors into these internal and external degrees of freedom, while the full G is indeed the full unitary group - the difference between those internal and external degrees of freedom disappears at high energies.

Technically, the unitary extensions correspond of course to self-adjoint extensions of the fuzzy X_i beyond their domain of definition (beyond e.g. where the stringy uncertainty relation holds). In terms of infinite dimensional matrices, one thereby adds to the domain of an X a maximal set of vectors for which the limits $n_1 \rightarrow \infty, n_2 \rightarrow \infty$ of Eq.1.3 commute. Intuitively, these vectors (because they have an infinite number of nonzero entries) explore the matrix X_{ij} at $i, j = \infty$.

Within any one self-adjoint extended domain the limits $n_1, n_2 \rightarrow \infty$ commute i.e., in the jargon of the first page, the vectors in this domain all probe X_{ij} at the same corner of the infinity $i, j = \infty$. The set of different self-adjoint domain extensions forms a representation of T , and each extension probes the behaviour of X_{ij} at different corners of the infinity $i, j = \infty$.

Thus, while the fuzzy X_i show only those degrees of freedom that correspond to structure larger than the cutoff scale $(\Delta X)_0$, the unitaries in G , being bounded operators, live on the entire Hilbert space and they therefore must ‘see’ all degrees of freedom. If a fundamental quantum gravity theory does involve fuzzy X_i , then it seems, therefore, that the degrees of freedom beyond the cutoff scale naturally reappear as internal degrees of freedom, with a gauge group structure related to T .

7. REMARKS ON THE POSSIBILITY OF SYMMETRIC OR ISOMETRIC OPERATORS OCCURRING IN LARGE SCALE GRAVITY

Finally, let us end with some speculations on the possibility of simple symmetric or isometric matrices appearing with quantum gravity effects at large scales.

We recall that the notion of a particle's energy-momentum is a delicate one. The fact that there exists a covariant definition of a phase space of fields and their conjugate *momentum fields* does not (since momentum *fields* act as translators in the space of field amplitudes rather than as translators on the space-time manifold) appear to resolve a fundamental paradox:

Classically, the momentum of a particle is the *local* tangent to its geodesic trajectory, while quantum theoretically a particle (as far as the notion of particle goes on curved space) which has a fairly certain momentum is highly *delocalised*. On flat space one can add and average the momenta on different virtual trajectories to obtain momentum expectations. But on an even only slightly curved space this summation must, at least, pick up uncertainties - because the tangent vectors live in different tangent spaces and parallel transport is then nonunique since path dependent. Thus, the more a space is curved, the more the notion of a particle's energy-momentum is imprecise.

One may therefore be tempted to give up on a notion of a particle's momentum on a curved space. After all, on flat space one is accustomed to the existence of plane waves which stand for the "points" that make up momentum space, while on a generic curved space there is no covariant notion of plane waves.

On the other hand, there is the Einstein equation: while the curvature on the left hand side is local, the energy and momentum on the right-hand side are, quantum mechanically, intrinsically nonlocal, a feature which may in some form persist to more fundamental, quantised successors of the Einstein equation. Maybe, one therefore ultimately cannot resolve the paradoxes of momenta on curved space by simply abandoning the concept of momenta.

Let us here only speculate that a notion of 'fuzzy' momentum operators as some form of translators on curved space may exist. We would expect them to be noncommutative to account for curvature, and we may expect them to be simple symmetric to account for the absence of plane waves, i.e. for the absence of points in momentum space. Finite translations in space or time could then be unitary or merely isometric. In particular, in the presence of horizons one may expect an isometric time evolution which does not have a unitary extension. An isometric time evolution operator would be able to preserve the scalar product on all fields where it is defined, without, however, being invertible. Some information may again be 'nontrivially stored', and as extension problems can always be mapped onto abstract boundary condition problems, this information may be viewed as sitting on the horizon. Technically, we would be dealing with the case of deficiency spaces of nonequal size. For example, the derivative $-i\partial_x$ on the half line is merely symmetric without self-adjoint extension. It gives rise to obviously noninvertible finite translations which are however still isometric i.e. scalar product preserving.

Properties of symmetric and isometric operators have of course long been known, see e.g. [10], but I hope to have shown that their potential for physical applications has not yet been fully explored.

I am grateful to the anonymous referee for pointing out a possible relation between the above picture which involves (nonresolvable) lattices at the Planck scale, and an old idea by Sakharov. In [11], Sakharov relates gravity and internal degrees of freedom at the Planck scale in order to motivate the Einstein-Hilbert action. His idea is that this action expresses a ‘metrical elasticity’ of space-time that arises from curvature-induced nonperfect matching of the zero-point energies in the vacuum of the QFT of the nongravitational (inner) degrees of freedom. See also [12]. The analogy suggested by Sakharov describes gravitons essentially as the phonons of the complicated vacuum of the internal degrees of freedom.

References

- [1] H.S. Snyder, Phys. Rev. 71: 38 (1947)
- [2] A. Connes, *Noncommutative Geometry*, AP (1994), G. Landi, *An Introduction to Noncommutative Spaces and their Geometry*, Lecture Notes in Physics: Monograph 51, Springer, Heidelberg (1997), J. Madore, *An introduction to noncommutative differential geometry and its physical applications*, CUP (1995), S. Majid, *Foundations of Quantum Group Theory*, CUP (1996)
- [3] S. Doplicher, K. Fredenhagen, J.E. Roberts, Comm. Math. Phys. **172**, 187 (1995), D.V. Ahluwalia, Phys. Lett. **B339**, 301 (1994), M.-J. Jaeckel, S. Reynaud, Phys. Lett. **A185**, 143 (1994), A. Connes, M.R. Douglas, A. Schwarz, RU-97-94, hep-th/9711162, C. Rovelli, Preprint C-97-12-16, gr-qc/9803024
- [4] E. Witten, Phys. Today **49** (4), 24 (1996), D.J. Gross, P.F. Mende, Nucl. Phys. **B303**, 407 (1988), D. Amati, M. Ciafaloni, G. Veneziano, Phys.Lett. **B216** 41, (1989), K. Konishi, G. Paffuti, P. Provero, Phys. Lett. **B234**, 276 (1990), R. Guida, K. Konishi, P. Provero, Mod. Phys. Lett. **A6**, 1487 (1991), L.J. Garay, Int. J. Mod. Phys. **A10**, 145 (1995)
- [5] A. Kempf, J. Phys. **A30** 2093, (1997)
- [6] A. Kempf, G. Mangano, Phys. Rev. **D55**, 7909 (1997)
- [7] A. Kempf, in Proc. 21ICGTMP Goslar, hep-th/9612082 (1997)
- [8] A. Kempf, Europhys. Lett. **40** (3), 257 (1997)
- [9] A. Kempf, in preparation
- [10] N.I. Akhiezer, I.M. Glazman, *Theory of Linear Operators in Hilbert space*, F. Unger Publ., (1963)
- [11] A.D. Sakharov, Soviet Physics - Doklady, **12** (11), 1040 (1968)
- [12] C.W. Misner, K.S. Thorne, J.A. Wheeler, *Gravitation*, p.285 Freeman & Co (1973)